

Resource Summary Report

Generated by [dkNET](#) on Sep 16, 2026

NTS1 antibody

RRID:AB_625999

Type: Antibody

Proper Citation

GeneTex Cat# GTX80747, RRID:AB_625999

Antibody Information

URL: http://antibodyregistry.org/AB_625999

Proper Citation: GeneTex Cat# GTX80747, RRID:AB_625999

Target Antigen: NTS1 antibody

Host Organism: guinea pig

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: Immunocytochemistry. Immunocytochemistry 1:1000 (Dilutions listed only as a recommendation. Optimal dilution should be determined by investigator.) Immunocytochemistry Transfected COS-7 cells, plated on poly-L-lysine-coated glass coverslips, were fixed for 20 minutes with 4 paraformaldehyde in 0.1 M phosphate buffer (PB), pH 7.4, rinsed with 0.1 M Tris base-buffered saline (TBS), pH 7.4, and preincubated for 30 minutes at RT with a blocking solution consisting of 5 normal goat serum (NGS), 2 BSA, and 0.1 Triton X-100 in 0.1 M TBS. Cells were then incubated overnight at 4°C with the anti-NTS1 peptide antiserum diluted 1: 1,000 in 0.1 M TBS, pH 7.4, containing 1 NGS and 0.05 Triton X-100. After washing with TBS, cells were incubated for 1 hour at RT with appropriate second

Antibody Name: NTS1 antibody

Description: This polyclonal targets NTS1 antibody

Target Organism: rat, others not tested

Antibody ID: AB_625999

Vendor: GeneTex

Catalog Number: GTX80747

Record Creation Time: 20231110T080400+0000

Ratings and Alerts

No rating or validation information has been found for NTS1 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunocytochemistry.

Immunocytochemistry 1:1000 (Dilutions listed only as a recommendation. Optimal dilution should be determined by investigator.) Immunocytochemistry Transfected COS-7 cells, plated on poly-L-lysine-coated glass coverslips, were fixed for 20 minutes with 4 paraformaldehyde in 0.1 M phosphate buffer (PB), pH 7.4, rinsed with 0.1 M Trisma base-buffered saline (TBS), pH 7.4, and preincubated for 30 minutes at RT with a blocking solution consisting of 5 normal goat serum (NGS), 2 BSA, and 0.1 Triton X-100 in 0.1 M TBS. Cells were then incubated overnight at 4°C with the anti-NTS1 peptide antiserum diluted 1: 1,000 in 0.1 M TBS, pH 7.4, containing 1 NGS and 0.05 Triton X-100. After washing with TBS, cells were incubated for 1 hour at RT with appropriate seconda

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.