

# Resource Summary Report

Generated by [dkNET](#) on Sep 4, 2026

## Donkey anti-Goat IgG (H+L) Secondary Antibody, FITC

RRID:AB\_2534674

Type: Antibody

---

### Proper Citation

Thermo Fisher Scientific Cat# A16000, RRID:AB\_2534674

---

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_2534674](http://antibodyregistry.org/AB_2534674)

**Proper Citation:** Thermo Fisher Scientific Cat# A16000, RRID:AB\_2534674

**Target Antigen:** Goat IgG (H+L)

**Host Organism:** donkey

**Clonality:** polyclonal secondary

**Comments:** Applications: ICC/IF (1:500-1:20,000), Flow (1:500-1:20,000)

**Antibody Name:** Donkey anti-Goat IgG (H+L) Secondary Antibody, FITC

**Description:** This polyclonal secondary targets Goat IgG (H+L)

**Target Organism:** goat

**Antibody ID:** AB\_2534674

**Vendor:** Thermo Fisher Scientific

**Catalog Number:** A16000

**Record Creation Time:** 20231110T035518+0000

**Record Last Update:** 20260830T013330+0000

---

### Ratings and Alerts

No rating or validation information has been found for Donkey anti-Goat IgG (H+L) Secondary Antibody, FITC.

No alerts have been found for Donkey anti-Goat IgG (H+L) Secondary Antibody, FITC.

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We found 3 mentions in open access literature.

**Listed below are recent publications.** The full list is available at [dkNET](#) .

Magno JC, et al. (2026) Engineering T Cells with a Tumor-Reactive Chimeric T-cell Receptor Reduces Exhaustion and Promotes Persistence to Elicit Enhanced Antitumor Responses. Cancer research, 86(11), 2688.

Hsieh LT, et al. (2025) The Mycobacterium ulcerans toxin mycolactone causes destructive Sec61-dependent loss of the endothelial glycocalyx and vessel basement membrane to drive skin necrosis. eLife, 12.

Li W, et al. (2023) Monocyte-derived Kupffer cells dominate in the Kupffer cell pool during liver injury. Cell reports, 42(10), 113164.