

# Resource Summary Report

Generated by [dkNET](#) on Aug 19, 2026

## CDK2 [2B6] antibody

RRID:AB\_369268

Type: Antibody

### Proper Citation

GeneTex Cat# GTX22363, RRID:AB\_369268

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_369268](http://antibodyregistry.org/AB_369268)

**Proper Citation:** GeneTex Cat# GTX22363, RRID:AB\_369268

**Target Antigen:** CDK2 [2B6] antibody

**Host Organism:** mouse

**Clonality:** monoclonal

**Comments:** Discontinued; manufacturer recommendations: IHC: This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 1 mM EDTA buffer, pH 8.0 prior to immunostaining., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, WB; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; Western Blot

**Antibody Name:** CDK2 [2B6] antibody

**Description:** This monoclonal targets CDK2 [2B6] antibody

**Target Organism:** rat, mouse, human

**Antibody ID:** AB\_369268

**Vendor:** GeneTex

**Catalog Number:** GTX22363

**Record Creation Time:** 20250819T231701+0000

## Ratings and Alerts

No rating or validation information has been found for CDK2 [2B6] antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: IHC: This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 1 mM EDTA buffer, pH 8.0 prior to immunostaining., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, WB; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; Western Blot

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We found 1 mentions in open access literature.

**Listed below are recent publications.** The full list is available at [dkNET](#) .

G??ttle P, et al. (2015) Oligodendroglial maturation is dependent on intracellular protein shuttling. The Journal of neuroscience : the official journal of the Society for Neuroscience, 35(3), 906.