

# Resource Summary Report

Generated by [dkNET](#) on Aug 16, 2026

## Insulin [K36aC10] antibody

RRID:AB\_375014

Type: Antibody

---

### Proper Citation

GeneTex Cat# GTX73586, RRID:AB\_375014

---

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_375014](http://antibodyregistry.org/AB_375014)

**Proper Citation:** GeneTex Cat# GTX73586, RRID:AB\_375014

**Target Antigen:** Insulin [K36aC10] antibody

**Host Organism:** mouse

**Clonality:** monoclonal

**Comments:** Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Suitable for formalin/paraffin tissues., The usefulness of this product in other applications has not been determined.

**Antibody Name:** Insulin [K36aC10] antibody

**Description:** This monoclonal targets Insulin [K36aC10] antibody

**Target Organism:** feline, rat, porcine, canine, rabbit, bovine, human, dog, sheep

**Antibody ID:** AB\_375014

**Vendor:** GeneTex

**Catalog Number:** GTX73586

**Record Creation Time:** 20250819T231007+0000

**Record Last Update:** 20250820T043005+0000

---

## Ratings and Alerts

No rating or validation information has been found for Insulin [K36aC10] antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Suitable for formalin/paraffin tissues., The usefulness of this product in other applications has not been determined.

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.