

# Resource Summary Report

Generated by [dkNET](#) on Aug 4, 2026

## pEntry Tshld-GFP

RRID:Addgene\_53211

Type: Plasmid

---

### Proper Citation

RRID:Addgene\_53211

---

### Plasmid Information

**URL:** <http://www.addgene.org/53211>

**Proper Citation:** RRID:Addgene\_53211

**Insert Name:** Tshld-GFP

**Bacterial Resistance:** Kanamycin

**Defining Citation:** [PMID:24615791](#)

**Vector Backbone Description:** Backbone Marker:invitrogen; Backbone Size:4470;  
Vector Backbone:pDONR 201; Vector Types;; Bacterial Resistance:Kanamycin

**Comments:** The Tshld system allows users to fine-tune the concentration of a protein of interest in mammalian cells by titrating the concentration of rapamycin or FK506. Unlike other post-translational techniques, the Tshld system releases its protein of interest in its native form, so it is highly attractive to be used in investigations of proteins involved in multimeric protein complexes.

**Plasmid Name:** pEntry Tshld-GFP

**Record Creation Time:** 20220422T222312+0000

**Record Last Update:** 20260801T081442+0000

---

### Ratings and Alerts

No rating or validation information has been found for pEntry Tshld-GFP.

No alerts have been found for pEntry Tshld-GFP.

---

### Data and Source Information

**Source:** [Addgene](#)

---

## Usage and Citation Metrics

We found 1 mentions in open access literature.

**Listed below are recent publications.** The full list is available at [dkNET](#) .

Gaynor AS, et al. (2017) Induced prodrug activation by conditional protein degradation. Journal of biotechnology, 260, 62.