

Resource Summary Report

Generated by [dkNET](#) on Jul 29, 2026

Japanese Ministry of Health Labor and Welfare

RRID:SCR_011326

Type: Tool

Proper Citation

Japanese Ministry of Health Labor and Welfare (RRID:SCR_011326)

Resource Information

URL: <http://www.mhlw.go.jp/english/>

Proper Citation: Japanese Ministry of Health Labor and Welfare (RRID:SCR_011326)

Description: Cabinet level ministry of the Japanese government. It is commonly known as K?r?-sh? in Japan. The ministry provides services on health, labour and welfare.

Abbreviations: Ministry of Health Labour and Welfare

Synonyms: Ministry of Health Labour and Welfare - Japan, Japanese Ministry of Health Labour and Welfare

Resource Type: institution

Keywords: Government granting agency

Resource Name: Japanese Ministry of Health Labor and Welfare

Resource ID: SCR_011326

Alternate IDs: Crossref funder ID: 501100003478, nlx_152042, Wikidata: Q1191238, grid.415828.2

Alternate URLs: <https://ror.org/03mwa7s65>

Record Creation Time: 20220129T080303+0000

Record Last Update: 20260725T190711+0000

Ratings and Alerts

No rating or validation information has been found for Japanese Ministry of Health Labor and Welfare.

No alerts have been found for Japanese Ministry of Health Labor and Welfare.

Data and Source Information

Source: [SciCrunch Registry](#)

Usage and Citation Metrics

We found 166 mentions in open access literature.

Listed below are recent publications. The full list is available at [dkNET](#) .

Kuroda D, et al. (2025) Optimizing the breadth of SARS-CoV-2-neutralizing antibodies in vivo and in silico. *Human vaccines & immunotherapeutics*, 21(1), 2526873.

Tateishi Y, et al. (2025) Functional genomics reveals strain-specific genetic requirements conferring hypoxic growth in *Mycobacterium intracellulare*. *eLife*, 13.

Matsuda H, et al. (2025) Fluctuating or progressive hearing loss in the middle-to-high frequencies may suggest the neurotologic endotype of perilymphatic fistula without antecedent traumatic events. *Frontiers in neurology*, 16, 1571379.

Sasaki A, et al. (2024) Prevalence of perilymphatic fistula in patients with sudden-onset sensorineural hearing loss as diagnosed by Cochlin-tomoprotein (CTP) biomarker detection: its association with age, hearing severity, and treatment outcomes. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*, 281(5), 2373.

Matsuda H, et al. (2020) Congenital Membranous Stapes Footplate Producing Episodic Pressure-Induced Perilymphatic Fistula Symptoms. *Frontiers in neurology*, 11, 585747.

Yasuda S, et al. (2018) Tumorigenicity-associated characteristics of human iPS cell lines. *PloS one*, 13(10), e0205022.

Nakatochi M, et al. (2018) Prediction model for pancreatic cancer risk in the general Japanese population. *PloS one*, 13(9), e0203386.

Yamada K, et al. (2018) LatY136F knock-in mouse model for human IgG4-related disease. *PloS one*, 13(6), e0198417.

Kato T, et al. (2018) Optimising first- and second-line treatment strategies for untreated major depressive disorder - the SUN?D study: a pragmatic, multi-centre, assessor-blinded randomised controlled trial. *BMC medicine*, 16(1), 103.

Kitamura K, et al. (2018) Flap endonuclease 1 is involved in cccDNA formation in the hepatitis B virus. *PLoS pathogens*, 14(6), e1007124.

Kikuchi H, et al. (2018) Short-term prognosis of emergently hospitalized dialysis-independent chronic kidney disease patients: A nationwide retrospective cohort study in Japan. *PloS one*, 13(11), e0208258.

Sato'o Y, et al. (2018) Tailor-made gene silencing of *Staphylococcus aureus* clinical isolates by CRISPR interference. *PloS one*, 13(1), e0185987.

Tsuboi M, et al. (2018) Clinical characteristics and epidemiology of intestinal tapeworm infections over the last decade in Tokyo, Japan: A retrospective review. *PLoS neglected tropical diseases*, 12(2), e0006297.

Cevallos W, et al. (2018) Diagnosis of amphimeriasis by LAMPhimerus assay in human stool samples long-term storage onto filter paper. *PloS one*, 13(2), e0192637.

Ikezono T, et al. (2018) The diagnostic performance of a novel ELISA for human CTP (Cochlin-tomoprotein) to detect perilymph leakage. *PloS one*, 13(1), e0191498.

Chida J, et al. (2018) Prion protein protects mice from lethal infection with influenza A viruses. *PLoS pathogens*, 14(5), e1007049.

Okuda K, et al. (2018) Comprehensive behavioral analysis of the Cdkl5 knockout mice revealed significant enhancement in anxiety- and fear-related behaviors and impairment in both acquisition and long-term retention of spatial reference memory. *PloS one*, 13(4), e0196587.

Eto S, et al. (2018) Mesenchymal stem cells derived from human iPS cells via mesoderm and neuroepithelium have different features and therapeutic potentials. *PloS one*, 13(7), e0200790.

Horibe Y, et al. (2017) Alpha-glucosidase inhibitor use is associated with decreased colorectal neoplasia risk in patients with type 2 diabetes mellitus receiving colonoscopy: a retrospective study. *Oncotarget*, 8(58), 97862.

Cevallos W, et al. (2017) LAMPhimerus: A novel LAMP assay for detecting *Amphimerus* sp. DNA in human stool samples. *PLoS neglected tropical diseases*, 11(6), e0005672.