## **Resource Summary Report**

Generated by dkNET on Apr 29, 2025

# **HMCan**

RRID:SCR\_010858

Type: Tool

## **Proper Citation**

HMCan (RRID:SCR\_010858)

#### **Resource Information**

URL: http://www.cbrc.kaust.edu.sa/hmcan/

Proper Citation: HMCan (RRID:SCR\_010858)

Description: A Hidden Markov Model based software tool that is developed to detect

histone modification in cancer ChIP-seq data.

**Abbreviations:** HMCan

**Synonyms:** Histone Modification in Cancer

**Resource Type:** software resource

**Defining Citation: PMID:24021381** 

Keywords: bio.tools

**Funding:** 

Resource Name: HMCan

Resource ID: SCR\_010858

Alternate IDs: biotools:hmcan, OMICS\_00443

Alternate URLs: https://bio.tools/hmcan

**Record Creation Time:** 20220129T080301+0000

**Record Last Update:** 20250420T014512+0000

### **Ratings and Alerts**

No rating or validation information has been found for HMCan.

No alerts have been found for HMCan.

#### Data and Source Information

Source: SciCrunch Registry

## **Usage and Citation Metrics**

We found 10 mentions in open access literature.

**Listed below are recent publications.** The full list is available at dkNET.

Thirant C, et al. (2023) Reversible transitions between noradrenergic and mesenchymal tumor identities define cell plasticity in neuroblastoma. Nature communications, 14(1), 2575.

Mohammed Ismail W, et al. (2023) MacroH2A histone variants modulate enhancer activity to repress oncogenic programs and cellular reprogramming. Communications biology, 6(1), 215.

Gregoricchio S, et al. (2022) HDAC1 and PRC2 mediate combinatorial control in SPI1/PU.1-dependent gene repression in murine erythroleukaemia. Nucleic acids research, 50(14), 7938.

Jdeed S, et al. (2022) The Role of ARID1A in the Nonestrogenic Modulation of IGF-1 Signaling. Molecular cancer research: MCR, 20(7), 1071.

Jarroux J, et al. (2021) HOTAIR IncRNA promotes epithelial-mesenchymal transition by redistributing LSD1 at regulatory chromatin regions. EMBO reports, 22(7), e50193.

Liehrmann A, et al. (2021) Increased peak detection accuracy in over-dispersed ChIP-seq data with supervised segmentation models. BMC bioinformatics, 22(1), 323.

Kaukonen D, et al. (2020) Analysis of H3K4me3 and H3K27me3 bivalent promotors in HER2+ breast cancer cell lines reveals variations depending on estrogen receptor status and significantly correlates with gene expression. BMC medical genomics, 13(1), 92.

Erd?s E, et al. (2020) NR2F2 Orphan Nuclear Receptor is Involved in Estrogen Receptor Alpha-Mediated Transcriptional Regulation in Luminal A Breast Cancer Cells. International journal of molecular sciences, 21(6).

Lopez-Delisle L, et al. (2018) Activated ALK signals through the ERK-ETV5-RET pathway to drive neuroblastoma oncogenesis. Oncogene, 37(11), 1417.

Ashoor H, et al. (2017) HMCan-diff: a method to detect changes in histone modifications in

cells with different genetic characteristics. Nucleic acids research, 45(8), e58.