

Resource Summary Report

Generated by [dkNET](#) on Sep 20, 2026

MojoSort™ Mouse NK Cell Isolation Kit Column Protocol

DOI:10.17504/protocols.io.7bhhij6

Type: Protocol

Proper Citation

Sam Li 2019. MojoSort™ Mouse NK Cell Isolation Kit Column Protocol. protocols.io dx.doi.org/10.17504/protocols.io.7bhhij6

Protocol Information

URL: <https://dx.doi.org/10.17504/protocols.io.7bhhij6>

Authors: Sam Li

Group: BioLegend

Summary: BioLegend MojoSort™ nanobeads work in commonly used separation columns, based on our internal research as well as validation by external testing by academic labs. This simple protocol consists of following the MojoSort™ protocol to label the cells with pre-diluted MojoSort™ reagents and using the columns as indicated by the manufacturer. Note: Due to the properties of our beads, it may be possible to use far fewer beads and less antibody cocktail than with other commercial suppliers. We recommend a titration to find the best dilution factor. However, as a general rule, dilutions ranging from 1:2 to 1:10 for the antibody cocktail can be used. Dilutions ranging from 1:5 to 1:20 for the Streptavidin Nanobeads can be used. Please contact BioLegend Technical Service (tech@biolegend.com) if further assistance is needed.

Affiliations: BioLegend

External URL: <https://www.biolegend.com/protocols/mojosort-mouse-nk-cell-isolation-kit-column-protocol/4771/>

Version: 1

Publication Date: 2019

Proper Citation: Sam Li 2019. MojoSort™ Mouse NK Cell Isolation Kit Column Protocol. protocols.io dx.doi.org/10.17504/protocols.io.7bhhij6

Record Creation Time: 20210329T220546+0000

Record Last Update: 20210329T221158+0000

Data and Source Information

Source: