

Resource Summary Report

Generated by [dkNET](#) on Sep 5, 2026

The ARF-AID system: Methods that preserve endogenous protein levels and facilitate rapidly inducible protein degradation

DOI:10.17504/protocols.io.bc7cziw

Type: Protocol

Proper Citation

Kizhakke Mattada Sathyan, Thomas G. Scott, Michael J. Guertin 2020. The ARF-AID system: Methods that preserve endogenous protein levels and facilitate rapidly inducible protein degradation. protocols.io dx.doi.org/10.17504/protocols.io.bc7cziw

Protocol Information

URL: <https://dx.doi.org/10.17504/protocols.io.bc7cziw>

Authors: Kizhakke Mattada Sathyan, Thomas G. Scott, Michael J. Guertin

Summary: The ARF-AID (Auxin Response Factor-Auxin Inducible Degron) system is a re-engineered auxin-inducible protein degradation system. Inducible degron systems are widely used to specifically and rapidly deplete proteins of interest in cell lines and organisms. An advantage of inducible degradation is that the biological system under study remains intact and functional until perturbation. This feature necessitates that the endogenous levels of the protein are maintained. However, endogenous tagging of genes with AID can result in chronic, auxin-independent proteasome-mediated degradation. The additional expression of the ARF-PB1 domain in the re-engineered ARF-AID system prevents chronic degradation of AID-tagged proteins while preserving rapid degradation of tagged proteins. Here we describe the protocol for engineering human cell lines to implement the ARF-AID system for specific and inducible protein degradation. These methods are adaptable and can be extended from cell lines to organisms.

Affiliations: Biochemistry and Molecular Genetics Department, Biochemistry and Molecular Genetics Department, Biochemistry and Molecular Genetics Department; Center for Public Health Genomics; Cancer Center; University of Virginia

Version: 1

Publication Date: 2020

Proper Citation: Kizhakke Mattada Sathyan, Thomas G. Scott, Michael J. Guertin 2020. The ARF-AID system: Methods that preserve endogenous protein levels and facilitate

rapidly inducible protein degradation. protocols.io
dx.doi.org/10.17504/protocols.io.bc7cziw

Record Creation Time: 20210329T220539+0000

Record Last Update: 20210329T221055+0000

Data and Source Information

Source: