

Resource Summary Report

Generated by [dkNET](#) on Sep 5, 2026

ELISA for quantification of IL-36 in human serum.

DOI:10.17504/protocols.io.bj38kqrw

Type: Protocol

Proper Citation

Angel Justiz-Vaillant 2020. ELISA for quantification of IL-36 in human serum.. protocols.io dx.doi.org/10.17504/protocols.io.bj38kqrw

Protocol Information

URL: <https://dx.doi.org/10.17504/protocols.io.bj38kqrw>

Authors: Angel Justiz-Vaillant

Group: University of the West Indies, angel.vaillant@sta.uwi.edu

Summary: Interleukins (IL) are a type of cytokine first thought to be expressed by leukocytes alone but have later been found to be produced by many other body cells. They play essential roles in the activation and differentiation of immune cells, as well as proliferation, maturation, migration, and adhesion. They also have pro-inflammatory and anti-inflammatory properties. The primary function of interleukins is, therefore, to modulate growth, differentiation, and activation during inflammatory and immune responses. Interleukins consist of a large group of proteins that can elicit many reactions in cells and tissues by binding to high-affinity receptors in cell surfaces. Phagocytes mainly make IL-36. It acts on T lymphocytes and NK cells regulating the IFN-?? synthesis. It stimulates the hematopoiesis and expression of both MHC class I and II molecules as well as intracellular adhesion molecules (ICAM)-1. [1]Reference1. Justiz Vaillant AA, Qurie A. Interleukin. In:StatPearls. Treasure Island (FL): StatPearls Publishing; June 12, 2019.

Affiliations: University of the West Indies St. Augustine

Version: 1

Publication Date: 2020

Proper Citation: Angel Justiz-Vaillant 2020. ELISA for quantification of IL-36 in human serum.. protocols.io dx.doi.org/10.17504/protocols.io.bj38kqrw

Record Creation Time: 20210329T220539+0000

Record Last Update: 20210329T221056+0000

Data and Source Information

Source: