

Resource Summary Report

Generated by [dkNET](#) on Sep 12, 2026

ELISA for quantification of human immunoglobulin D (IgD) in serum or plasma.

DOI:10.17504/protocols.io.bj7wkrpe

Type: Protocol

Proper Citation

Angel Justiz-Vaillant, Belkis Ferrer-Cosme 2020. ELISA for quantification of human immunoglobulin D (IgD) in serum or plasma.. protocols.io
dx.doi.org/10.17504/protocols.io.bj7wkrpe

Protocol Information

URL: <https://dx.doi.org/10.17504/protocols.io.bj7wkrpe>

Authors: Angel Justiz-Vaillant, Belkis Ferrer-Cosme

Group: University of the West Indies, angel.vaillant@sta.uwi.edu

Summary: IgD is a monomer with a molecular weight of 184 Kd. IgD is present in a meager amount in the serum (0.03 mg/mL) and has an unknown function against pathogens. It is regarded as a BCR. IgD may play an essential role in antigen-triggered lymphocyte differentiation. [1]

Affiliations: University of the West Indies St. Augustine, "Saturnino Lora Torres" Provincial Teaching Clinical Surgical Hospital. Cuba

Version: 1

Publication Date: 2020

Proper Citation: Angel Justiz-Vaillant, Belkis Ferrer-Cosme 2020. ELISA for quantification of human immunoglobulin D (IgD) in serum or plasma.. protocols.io
dx.doi.org/10.17504/protocols.io.bj7wkrpe

Record Creation Time: 20210329T220519+0000

Record Last Update: 20210329T220843+0000

Data and Source Information

Source: