

Resource Summary Report

Generated by [dkNET](#) on Sep 13, 2026

Laboratory protocols for the study “Whole genome analysis of selected human and animal rotaviruses identified in Uganda from 2012 to 2014 reveals complex genome re-assortment events between human, bovine, caprine and porcine strains”.

DOI:10.17504/protocols.io.h4vb8w6

Type: Protocol

Proper Citation

Josephine Bwogi^{1, 2}, Khuzwayo C. Jere^{3, 4}, Charles Karamagi², Denis K. Byarugaba⁵, Prossy Namuwulya¹, Frederick N. Baliraine⁶, Ulrich Desselberger⁷, Miren Iturriza-Gomara³. 2017. Laboratory protocols for the study “Whole genome analysis of selected human and animal rotaviruses identified in Uganda from 2012 to 2014 reveals complex genome re-assortment events between human, bovine, caprine and porcine strains”.. protocols.io dx.doi.org/10.17504/protocols.io.h4vb8w6

Protocol Information

URL: <https://dx.doi.org/10.17504/protocols.io.h4vb8w6>

Authors: Josephine Bwogi^{1, 2}, Khuzwayo C. Jere^{3, 4}, Charles Karamagi², Denis K. Byarugaba⁵, Prossy Namuwulya¹, Frederick N. Baliraine⁶, Ulrich Desselberger⁷, Miren Iturriza-Gomara³.

Associated Publications: Bwogi J, Jere KC, Karamagi C, Byarugaba DK, Namuwulya P, Baliraine FN, Desselberger U, Iturriza-Gomara M (2017) Whole genome analysis of selected human and animal rotaviruses identified in Uganda from 2012 to 2014 reveals complex genome reassortment events between human, bovine, caprine and porcine strains. PLoS ONE 12(6): e0178855. doi: [10.1371/journal.pone.0178855](https://doi.org/10.1371/journal.pone.0178855)

Affiliations: 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda

6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA
7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda
6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA
7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda
6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA
7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda
6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA
7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda
6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA
7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding

author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda 6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA 7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda 6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA 7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda 6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA 7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug, 1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda 6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA 7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug,

1 EPI laboratory, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda 2Department of Paediatrics and Child Health, College of Health Sciences, Makerere University, P.O. BOX 7072, Kampala Uganda 3 Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Liverpool, UK 4 Malawi-Liverpool-Wellcome Trust Clinical Research Programme / Department of Medical Laboratory Sciences, University of Malawi, College of Medicine, Blantyre, Malawi 5 Department of Microbiology, College of Veterinary Medicine and Biosecurity, Makerere University, P.O.BOX 7072, Kampala, Uganda 6Department of Biology and Kinesiology, LeTourneau University, Longview, Texas, USA 7Department of Medicine, University of Cambridge, Cambridge, UK *Corresponding author: Josephine Bwogi, Uganda Virus Research Institute, 51-59 Nakiwogo Road, P.O.BOX 49, Entebbe, Uganda. Email: josephinebwn@yahoo.co.uk or jbwogi@uvri.go.ug

External URL: <https://doi.org/10.1371/journal.pone.0178855>

Version: 1

Publication Date: 2017

Proper Citation: Josephine Bwogi^{1, 2}, Khuzwayo C. Jere^{3, 4}, Charles Karamagi², Denis K. Byarugaba⁵, Prossy Namuwulya¹, Frederick N. Baliraine⁶, Ulrich Desselberger⁷, Miren Iturriza-Gomara³. 2017. Laboratory protocols for the study “Whole genome analysis of selected human and animal rotaviruses identified in Uganda from 2012 to 2014 reveals complex genome re-assortment events between human, bovine, caprine and porcine strains”.. protocols.io dx.doi.org/10.17504/protocols.io.h4vb8w6

Record Creation Time: 20210329T220521+0000

Record Last Update: 20210329T220848+0000

Data and Source Information

Source: